ISU-IAP.TH 99-01,Irkutsk.

arXiv:hep-ph/9904300v1 12 Apr 1999

The polarization of a final electron in the field of intensive electromagnetic wave

E.M.Bol'shedvorsky and S.I.Polityko
Irkutsk State University
E-mail:polityko@psi.isu.ru

1999

Abstract

The polarization properties of the Compton effect in the field of a circularly polarized electromagnetic wave are discussed. In the case of nonlinear Compton effect the behavior of spin is analyzed. The transformation of the longitudinal polarization in the transverse polarization is showed. We obtained the total formulas describing the polarization of the final electron in the nonlinear case.

1.Introduction

The Compton effect is the main process obtaining from intensive high energy γ bunches on the base of the present linear electron collider, so named electron-photon and photon-photon collideres (PLC) [1],[2]. The polarization properties of this process are very important. For example, if the helities of initial particles are opposite the high energy part of spectrum increased twice comparing with the nonpolarizing case.

The optimal choice of the conversion coefficient is important in the conversion scheme (the relation between hard photon number to number of electron in bunches). The conversion coefficient is proportional to the laser flash energy and its increasing in the conversion region causes the important processes of multiple scattering electrons on a laser photon . In this case the effects of nonlinear quantum electrodinamics are essential [3]. In this study we consider the polarization of final electron in polarized electron scattering on a circularly polarized laser photons via absorption of s photons from a laser wave and perform summation over the final-photon polarizations. Some particular cases in the problem of final electron polarization in nonlinear QED were treated in [4] but the leading formulas can be used only in the c.m.s. of colliding particles. However, $e \to \gamma$ - conversion must be considered in a different frame where an electron with energy $E \sim 100$ collides with a laser photon with energy $\omega \sim 1$. This frame will be referred as a collider frame. The linear case of this problem was considered earlier in [5].

We shall consider the process of photon emission by an electron via simultaneous absorption of photons from a laser wave (nonlinear Compton effect).

$$s\gamma_0 + e^- \to \gamma + e^- \tag{1}$$

The systematic investigation of such a process was performed in [7],[8]. The possibility of observing nonlinear QED effects for an unpolarized electron beam in reaction (1) was discussed earlier in [3]. It was shown in [2],[6] that the application of longitudinally polarized electron beams and circularly polarized laser light leads to substantial improvements in the monochromaticity of $\gamma\gamma$ collisions. Anyway probability, there are the initial-beam polarizations that will be used in the implementation of concepts based on colliding γe and $\gamma\gamma$ beams.

However, when we investigate the Compton scattering of a high energy electron by a laser beam ($e \to \gamma$ conversion) at a comparatively large conversion ratio, we must take into consideration multiple electron scattering by laser photons. Because the required formulas were not available the similar calculations have already been performed in papers [9], but they either considered the conversion of unpolarized particles or disregarded the change in electron polarization in the scattering process . However, it is necessary to know the polarization of the final electron, since it is the initial electron for the next scattering event.

In this paper we calculate the polarization of the final electron and discuss its characteristics features. It is found that polarization effects are essential and that they must be taken into account in calculations dealing with multiple scattering.

2. The bases for polarization describing of initial and final particles

Nonlinear QED processes are characterized by the parameter

$$\xi^2 = \frac{e^2 F^2 \hbar^2}{m^2 \omega_0^2 c^2},$$

where F-is the strength of the electromagnetic-wave field, m-is the mass of electron, ω_0 is the energy of laser photon. At $\xi^2 \ll 1$ an electron interacts only with one photon (usual Compton effect), at $\xi^2 \gg 1$ an electron interacts with a collective field (synchrotron radiation). The transverse motion of electron in the electromagnetic-wave field leads to increase the effective electron mass $m_*^2 = m^2(1+\xi^2)$ and changes the spectrum of scattering photons: the high energy part of the spectrum with $s = 1$ squeezes and the high energy tail appears, corresponding to the absorption from the wave two or more laser photons. Considering that the electron in the electromagnetic-wave field becomes "heavier" we can write the energy-momentum conservation law in the form

$$q_1 + s k_1 = q_2 + k_2,$$

where q_1, q_2 "quasimomentums" of initial and final electrons

$$q_1 = p_1 + \xi^2 \frac{m^2}{2(k_1 p_1)} k_1, \qquad q_2 = p_2 + \xi^2 \frac{m^2}{2(k_1 p_2)} k_1.$$

Let us introduce the invariants for describing of scattering process:

$$x = \frac{2 p_1 k_1}{m^2}, \quad y = \frac{k_1 k_2}{p_1 k_1}.$$

Let $\boldsymbol{\zeta}$ and $\boldsymbol{\zeta'}$ — be the polarization vectors of the initial and final electrons. They determine the electron-spin 4-vectors as

$$\sigma = \left(\frac{\boldsymbol{\zeta}\mathbf{p_1}}{m}\,, \boldsymbol{\zeta} + \mathbf{p_1}\frac{\boldsymbol{\zeta}\mathbf{p_1}}{m(E_1+m)} \right)\,, \quad \sigma' = \left(\frac{\boldsymbol{\zeta'}\mathbf{p_2}}{m}\,, \boldsymbol{\zeta'} + \mathbf{p_2}\frac{\boldsymbol{\zeta'}\mathbf{p_2}}{m(E_2+m)} \right)$$

and the average helicities of the initial and final electrons as

$$\lambda_e = \frac{\boldsymbol{\zeta}\mathbf{p_1}}{2|\mathbf{p_1}|}, \quad \lambda_e' = \frac{\boldsymbol{\zeta}'\mathbf{p_2}}{2|\mathbf{p_2}|}.$$

In the collider system we choose the z axis along the initial electron momentum. We denote by θ and φ the polar and azimuthal angles of final photon emission. All azimuthal angles are defined with respect to the fixed x axis. In this frame, the expression to which the invariants are reduced for $\theta \ll 1$ are given by

$$x = \frac{4E_1\omega_1}{m^2}, \quad y = \frac{\omega_2}{E} = 1 - \frac{E_2}{E_1};$$

$$\zeta_1 = \boldsymbol{\zeta}\,\frac{[\mathbf{k_1}, \mathbf{k_2}]}{|[\mathbf{k_1}, \mathbf{k_2}]|} = \zeta_\perp \sin(\varphi - \beta),$$

$$\zeta_2 = \boldsymbol{\zeta}\,\frac{\mathbf{k_{2\perp}}}{|\mathbf{k_{2\perp}}|} = \zeta_\perp \cos(\varphi - \beta), \quad \zeta_3 = 2\lambda_e,$$

where ω_2 and E_2 are the energies of the final photon and electron, $\mathbf{k_{2\perp}}$ and $\boldsymbol{\zeta}_\perp$ are the transverse components of the vectors $\mathbf{k_2}$ and $\boldsymbol{\zeta}$ with respect to the vector $\mathbf{q}$, φ and β are the azimuthal angles of $\mathbf{k_2}$ and $\boldsymbol{\zeta}$; $\zeta_\perp = |\boldsymbol{\zeta}_\perp|$.

Thus, ζ_1 is the initial-electron transverse polarization that is orthogonal to the scattering plane, ζ_2 is the transverse polarization lying in this plane.

We decompose the spin 4-vectors on the orthogonal system of bases vectors The bases vectors, describing the polarization state of initial particles are given

$$e_0 = \frac{p_1}{m}, \quad e_1 = n_3,$$

$$e_2 = -\frac{z}{2\xi\sqrt{sxy}}x(1-y)(n_0 - n_1) - n_2, \quad e_3 = \frac{1}{m}(p_1 - \frac{2}{x}k).$$

For the final particles the basic vectors have the form

$$e_0' = \frac{p_2}{m}, \quad e_1' = n_3,$$

$$e_2' = -\frac{z}{2\xi\sqrt{sxy}}x(n_0 - n_1) - n_2, \quad e_3' = \frac{1}{m}(p_2 - \frac{2}{x(1-y)}k).$$

In this expressions

$$n_0 = \frac{K}{m\sqrt{sxy}}, \quad n_1 = \frac{q}{m\sqrt{sxy}}, \quad n_2 = \frac{\xi y}{mzx(1-y)}P\perp,$$

$$n_3^\alpha = \frac{\xi}{sm^3zx^2(1-y)}e^{\alpha\mu\rho\nu}K_\mu q_\rho P_\nu,$$

where

$$K = sk_1 + k_2, \quad q = k_2 - sk_1 = q_1 - q_2, \quad P\perp = P - \frac{PK}{K^2}K, \quad P = q_1 + q_2,$$

$$z = \frac{2s\xi}{\sqrt{1+\xi^2}}\sqrt{r(1-r)}, \quad r = \frac{y(1+\xi^2)}{(1-y)sx}.$$

These bases differ from the bases chosen in [5], here we have made a turn around the axes z' at the angle θ^* which is considered to be a scattering angle of a photon in the rest frame of initial electron

$$\cos\theta^* = 1 - \frac{2y}{sx(1-y) - \xi^2 y}.$$

We can represent the 4-vectors of spin σ and σ' as

$$\sigma=\sum_j \zeta_j e_j\ ,\quad \zeta_j=-\sigma e_j,\ ,\qquad\qquad \sigma'=\sum_j z'_j e'_j\ ,\quad z'_j=-\sigma' e'_j\,.$$

In the collider frame ζ' mean

$$\zeta'_1=\zeta'_\perp\sin(\varphi-\beta'),\quad \zeta'_2=\zeta'_\perp\cos(\varphi-\beta'),\quad \zeta'_3=2\lambda'_e,$$

where $\zeta'_\perp$ is the transverse component of the vector ζ' with respect to the vector $\mathbf{p_2}$, and β' is the azimuthal angle for the direction ζ'. Thus ζ'_3 is the double helicity of final electron, ζ'_2 is the transverse polarization lying in scattering plane, and ζ'_1 is orthogonal to this plane.The invariants ζ_j and z'_j completely characterize the polarization properties of electrons.

3. Differential probability of emission

We will calculate the probability of the process (1) using the method of quantum transition.As for the basis we use the exact nonstationary solutions of the Dirac equation (Volkov's solutions). We take into account the electron interaction with the external field . Electron interaction with the field of emitted photons is considered with the help of perturbation theory.

The amplitude of the electron transition from the state ψ_{p_1} in the wave field to the state ψ_{p_2} with the emission of a photon of momentum k_2 and polarization ε_2 is given by

$$M=e\int\bar\Psi_{p_2}(x)\hat\varepsilon_2^*\Psi_{p_1}(x)\frac{e^{-ik_2x}}{\sqrt{2E_2}}d^4x. \qquad (2)$$

where ε_2 is the final-photon polarization, $\Psi_{p_1}(x)$ $(\bar\Psi_{p_2}(x))$ is the exact solution of the Dirac equation for an electron in the field of a circularly polarized wave :

$$\Psi_{p_1}(x)=(1+\frac{ek_1\hat A}{2k_1p_1})u_{p_1}\exp(ie\frac{a_1p_1}{k_1p_1}\sin(\varphi)-ie\frac{a_2p_1}{k_1p_1}\cos(\varphi)+iqx),\quad \varphi=k_1x,$$

The vector potential of a circularly polarized electromagnetic wave has the form

$$A=a_1\cos\varphi+P_ca_2\sin\varphi,$$

where $k_1a_1=k_1a_2=a_1a_2=0,\quad a_1^2=a_2^2=a^2$. The average helicity of final (laser) photon P_c coincide with the Stokes parameter ξ_2. The density matrices of the polarized electrons are given by the standard form

$$\rho=\frac{1}{2}(\hat p_1+1)(1-\gamma^5\hat\sigma),\quad \rho'=\frac{1}{2}(\hat p_2+1)(1-\gamma^5\hat\sigma'),$$

where σ,σ' are 4-vectors of electron spin.

The density matrice of a final photon is

$$\nu_{ik}=e'_ie'^*_k=-\frac{g_{ik}}{2}$$

Using the standard techniques developed in [7],[8] we can easily calculate the probability of process (1) and perform summation on final-photon polarizations.

$$dW_s=\frac{\alpha m^2}{2E_1}\xi^2d\varphi dy\Big(F_{00}^{00}+P_c\sum_{i=1}^3(F_{i2}^{00}\zeta_i+F_{02}^{i0}\zeta'_i)+\sum_{i=1}^3\sum_{j=1}^3F_{i0}^{j0}\zeta_i\zeta'_j\Big), \qquad (3)$$

where $\alpha = e^2/4\pi$. Nonzero coefficients F_{ik}^{j0} are given

$$
\begin{aligned}
F_{00}^{00} &= -\frac{2}{\xi^2}J_s^2(z) + \Big(1 + \frac{y^2}{2(1-y)}\Big)(J_{s+1}^2(z) + J_{s-1}^2(z) - 2J_s^2(z)) \\
F_{32}^{00} = F_{02}^{30} &= J_s(z)J_s'(z)\frac{s(2-y)}{z}\Big(\frac{z^2x}{\xi^2 s} - 2\frac{y}{1-y}\Big) \\
F_{22}^{00} &= -2J_s(z)J_s'(z)\frac{y}{\xi} \\
F_{02}^{20} &= -2J_s(z)J_s'(z)\frac{y}{\xi(1-y)} \qquad (4) \\
F_{30}^{30} &= J_s^2(z)\frac{1}{\xi^2}\Big(2 - (2+\xi^2)\frac{y^2-2y+2}{1-y}\Big) + \frac{1}{2}\frac{y^2-2y+2}{1-y}(J_{s+1}^2(z) + J_{s-1}^2(z)) \\
F_{20}^{30} &= \Big(-\frac{2s}{z}\frac{y}{\xi} + \frac{z}{\xi^3}x(1-y)\Big)J_s^2(z) \\
F_{30}^{20} &= \Big(\frac{2s}{z}\frac{y}{\xi(1-y)} - \frac{z}{\xi^3}x\Big)J_s^2(z) \\
F_{20}^{20} = F_{10}^{10} &= \Big(J_{s+1}^2(z) + J_{s-1}^2(z)\Big) - 2\Big(\frac{1}{\xi^2} + 1\Big)J_s^2(z),
\end{aligned}
$$

where $J_s(z)$ is Bessel function.

4. Numerical calculations

For numerical calculations the physical parameters characterizing the region of laser conversion are chosen in accordance with [10]. In the conversion scheme it has been proposed to use a solid-state laser with laser photon energy $\omega_0 = 1.17$. Let us choose the parameter $x = 4.8$. Which is according to the threshold of e^+e^- pair production at collision of laser photon with high energy scattering photon. The length of the conversion region characterized by a high density of laser photon l_γ is close to the length of the electron bunch l_e. It is convenient to express the parameter ξ^2 in terms of the energy A, the duration τ and the radius a_γ of the laser beam at the point of intersection

$$\xi^2 = \frac{A}{A_*}, \quad \text{where} \quad A_* = \frac{\tau c}{4}\cdot\Big(\frac{m_e\omega_0 a_\gamma c}{e\hbar}\Big)^2.$$

The conversion coefficient k connect with the laser flash energy by relation

$$k = N_\gamma/N_e \sim 1 - \exp(-A/A_*) \sim A/A_*,$$

thus at $x = 4.8$ and $k = 1$ we get $\xi^2 = 0.05E[TeV]/l_\gamma$[10]. If $l_\gamma = 200\mu m$ and $E = 100$ Gev, then $\xi^2 = 0.25$. The expression (3) for the differential probability can be written in the following way

$$dW_s = \frac{\alpha m^2}{2E_1}\xi^2 d\varphi dy\Big(G_0 + \sum_{i=1}^{3} G_i\zeta_i'\Big),$$

where

$$
\begin{aligned}
G_0 &= F_{00}^{00} + P_cF_{32}^{00}\zeta_3 + P_cF_{22}^{00}\zeta_2, \\
G_1 &= F_{10}^{10}\zeta_1, \\
G_2 &= P_cF_{02}^{20} + F_{30}^{20}\zeta_3 + F_{20}^{20}\zeta_2, \\
G_3 &= P_cF_{02}^{30} + F_{30}^{30}\zeta_3 + F_{20}^{30}\zeta_2.
\end{aligned}
$$

The components of the polarization vector of final electron are given

$$\zeta_1'^f = \frac{G_1}{G_0}, \quad \zeta_2'^f = \frac{G_2}{G_0}, \quad \zeta_3'^f = \frac{G_3}{G_0}. \tag{5}$$

These formulas determine completely the polarization of final electron in the field of intensive electromagnetic wave. The case $\xi \gg 1$ is described in the Appendix.

In figures 1-6 the numerical calculations present the components of the polarization vector of final electron dependence $\zeta_i'^{(f)}$ as functions of its energy $y_1 = E'/E = 1 - y$ for various degrees of polarization of the initial electron and photon at $x = 4.8$. The number of absorbed laser photons $s = 5$ and the parameter $\xi^2 = 0.25$. Here $\zeta_1'^{(f)}$ is the final electron transverse polarization in the direction orthogonal to the scattering plane, $\zeta_2'^{(f)}$ is the transverse polarization in this plane and $\zeta_3'^{(f)}$ is the double average helicity. The analogous quantities for the initial electron are denoted by $\zeta_1, \ \zeta_2 \ \ \zeta_3 = 2\lambda_e$; P_c is the average helicity of initial photon. For the figures 1-6 we indicate nonzero components of the initial particles, for the figures 1-5 $\zeta_1'^{(f)} = 0$.

The appearance of the essential transverse polarization on the scattered electron is the common property shown in the figures. The transverse polarization takes place even in the case when the initial electron is not polarized. The electron polarization in the scattering plane changes the sign by altering the laser photon helicity (see.fig.1b and 2b). Hard scattering electrons for which $y_1 \approx 1$ conserve their helicity. In the soft region (small y_1) scattered electrons get the longitudional polarizations its sign depends on laser photon polarization sign. In this region the multiple photon absorption is essential.

We note, that at $\xi^2 \ll 1$ and $s = 1$ we get the results of[5].

We are very grateful to I. Ginzburg, G. Kotkin and V. Serbo for useful discussions.

Appendix

In the Appendix we shall consider the formulas for the probability of photon radiation by electron at large values ξ. The large value ξ can be obtained from decreasing the frequency at fixed field strength.

With increasing ξ the energy of wave the quant decreases but the number of absorption photons from electromagnetic wave increases $s \sim \xi^3$ and the mean energy and momentum of final electron increase effectively. The values are

$$x^* = \frac{2k_1p_1}{m^2}\xi, \quad y, \quad \rho_1 = \frac{\alpha_1}{8\beta}, \quad \rho_2 = \frac{\alpha_2}{8\beta},$$

where

$$\beta = -\frac{\xi^2 m^2}{8}\Big(\frac{1}{k_1p_1} - \frac{1}{k_1p_2}\Big) = \frac{\xi^2 y}{4x^*(1-y)},$$

does not depend on ξ. The variables $\rho_{1,2}$ are connected with azimuthal angle φ_0 as

$$\rho_1 = \frac{z}{8\beta}\cos\varphi_0, \quad \rho_2 = \frac{z}{8\beta}\sin\varphi_0,$$

where

$$\cos\varphi_0 = \frac{\alpha_1}{z}, \quad \sin\varphi_0 = \frac{\alpha_2}{z}.$$

Note that at $z \sim \xi^3$ the value $z/8\beta$ leads to unit effectively

We describe the difference of $z/8\beta$ from the unit as the new variable τ:

$$z = 8\beta\sqrt{1 + \frac{2\tau}{\xi}} = \frac{2\xi^3 y}{x^*(1-y)}\sqrt{1 + \frac{2\tau}{\xi}}.$$

Let's consider the probability of photon radiation by electron. Changing the order of integration and summation in the expression of the probability, we get

$$W=\sum_{s=1}^{\infty}\int_0^{2\pi}d\varphi\int_0^1 dyw=\int_0^{2\pi}d\varphi\int_0^1 dy\sum_{s>s_{min}}^{\infty}w,$$

where

$$s_{min}=\frac{\xi^3 y}{x^*(1-y)}\left(1+\frac{1}{\xi^2}\right).$$

At $\xi\gg 1$ the large number of terms and the sum can be changed by the integral. From the dependence z and$s$ it follows that

$$s=\frac{\xi^3 y}{x^*(1-y)}\left(1+\frac{2\tau}{\xi}\right)+s_{min},$$

Then

$$W=\int_0^{2\pi}d\varphi\int_0^1 dy\int_{-\xi^2/2}^{\infty}d\tau\frac{2\xi^2 y}{x^*(1-y)}w.$$

At $\xi\gg 1$ the value $z\sim s\sim\xi^3$, but their relation leads to the unit,

$$\sigma=\xi^2\left(1-\frac{z^2}{s^2}\right)=1+\tau^2,$$

and the function Bessel can be changed by the asymptotic Vatson representation

$$J_s(z)\approx\frac{1}{\pi}\left(\frac{2}{s}\right)^{1/3}Ai(\theta),\quad \theta=\left(\frac{s}{2}\right)^{2/3}\left(1-\frac{z^2}{s^2}\right)=\left(\frac{y}{x^*(1-y)}\right)^{2/3},$$

where $Ai(\theta)$ is the Eiri function.

Using this representation we finally get

$$W(\infty,x^*)=2\frac{\alpha m^2}{E_1\pi^2}\int_0^{2\pi}d\varphi\int_0^1 dy\int_{-\infty}^{\infty}d\tau\left(\frac{\theta}{\sigma}\right)^{1/2}\left(F_{00}^{00}+P_c\sum_{i=1}^{3}\left(F_{i2}^{00}\zeta_i+F_{02}^{i0}\zeta_i'\right)+\sum_{i=1}^{3}\sum_{j=1}^{3}F_{i0}^{j0}\zeta_i\zeta_j'\right),$$

where

$$F_{00}^{00}=-Ai^2(\theta)+\frac{\sigma}{\theta}(1+\frac{y^2}{2(1-y)})\left(\theta Ai^2(\theta)+\acute{A}i^2(\theta)\right)$$

$$F_{32}^{00}=F_{02}^{30}=-\left(\frac{\sigma}{\theta}\right)^{1/2}\frac{y(2-y)}{1-y}Ai(\theta)\acute{A}i(\theta)\tau$$

$$F_{22}^{00}=\left(\frac{\sigma}{\theta}\right)^{1/2}Ai(\theta)\acute{A}i(\theta)y$$

$$F_{02}^{20}=\left(\frac{\sigma}{\theta}\right)^{1/2}Ai(\theta)\acute{A}i(\theta)\frac{y}{1-y}$$

$$F_{30}^{30}=\frac{y-y^2-1}{1-y}Ai^2(\theta)+\frac{1}{2}\frac{y^2-2y+2}{1-y}\frac{\sigma}{\theta}\left(\theta Ai^2(\theta)+\acute{A}i^2(\theta)\right)$$

$$F_{20}^{30}=Ai^2(\theta)y\tau$$

$$F_{30}^{20}=Ai^2(\theta)\frac{y}{1-y}\tau$$

$$F_{20}^{20}=F_{10}^{10}=-Ai^2(\theta)+\frac{\sigma}{\theta}\left(\theta Ai^2(\theta)+\acute{A}i^2(\theta)\right)$$

The polarization of final electron is determined by formulae (5), where

$$G_0 = \int_{-\infty}^{\infty} d\tau \left(F_{00}^{00} + P_c F_{32}^{00} \zeta_3 + P_c F_{22}^{00} \zeta_2 \right),$$

$$G_1 = \int_{-\infty}^{\infty} d\tau \left(F_{10}^{10} \zeta_1 \right),$$

$$G_2 = \int_{-\infty}^{\infty} d\tau \left(P_c F_{02}^{20} + F_{30}^{20} \zeta_3 + F_{20}^{20} \zeta_2 \right),$$

$$G_3 = \int_{-\infty}^{\infty} d\tau \left(P_c F_{02}^{03} + F_{30}^{30} \zeta_3 + F_{20}^{30} \zeta_2 \right).$$

References

[1] *I.Ginzburg, G.Kotkin, V.Serbo, V.Telnov* Pizma ZhETF, 1981,**4**, 514;
Nucl. Instr. and Methods (NIM) 1983 **205**, 47;
I.F. Ginzburg, G.L. Kotkin, S.L. Panfil, V.G. Serbo, V.I. Telnov NIM, 1984 **219**, 5.

[2] *V.I.Telnov* Nucl.Instr. and Methods **24** (1990),72
I.F.Ginzburg Nucl.Instr. and Methods **A 355** (1995), 63

[3] *V.N.Baier, K.Yokoya* Particle Accelerators, 1994, Vol.44,pp.77-106
I.F.Ginzburg, G.L.Kotkin, S.I.Polityko Sov. Yad. Fiz., 1984, .40, 1495
I.F.Ginzburg, G.L.Kotkin, S.I.Polityko Sov. Yad. Fiz., 1993, .56, 65

[4] *K.A.Dyrgallo, S.I.Polityko* Phys. of Atomic Nuclei, 1995, v.58, 839

[5] *G.L,Kotkin, S.I.Polityko, V.G.Serbo* Phys. of Atomic Nuclei, 1996, v.59, 2147
G.L.Kotkin, S.I.Polityko, V.G.Serbo Nucl.Instr.and Methods **A 405** (1998),30

[6] *I.F.Ginzburg, G.L.Kotkin, S.L.Panfil, V.G.Serbo, V.I.Telnov* Sov. Yad. Fiz, v.38, 1021.

[7] *V.I.Ritus* Trudy FIAN,v.111,1979 [in Russian].

[8] *V.Berestetskii, E.Lifshitz and L.Pitaevskii* Quantum Electrodinamics, Pergamont press, Oxford, 1982.

[9] *Borden D.L., Bauer D.A., Caldwell D.O.* //Preprint SLAC-PUB-5715, Stanford, 1992;
Ginzburg I.F., Kotkin G.L., Polityko S.I. // Proc. Workshop on Physics at VLEPP, Moscow, 1992, p.27;

[10] *Telnov V.* Talk at ICFA Workshop *Quantum Aspect of Beam Physics*, Monterey, CA, USA, January 4-9,1998; hep-ex/9805002.

Figure 1: Components of the polarization vector of the scattered electron of its energy $\zeta_2'^{(f)}$, $\zeta_3'^{(f)}$ at $2\lambda_e = 1,\ P_c = -1$;

Figure 2: Components of the polarization vector of the scattered electron of its energy $\zeta_2'^{(f)}$, $\zeta_3'^{(f)}$ at $2\lambda_e = 1,\ P_c = +1$;

Figure 3: Components of the polarization vector of the scattered electron of its energy $\zeta_2'^{(f)}$, $\zeta_3'^{(f)}$ at $2\lambda_e = 0,\ P_c = +1$;

Figure 4: Components of the polarization vector of the scattered electron of its energy $\zeta_2^{\prime(f)}$, $\zeta_3^{\prime(f)}$ at $\zeta_2 = -1,\ P_c = 1$;

Figure 5: Components of the polarization vector of the scattered electron of its energy $\zeta_2^{\prime(f)}$, $\zeta_3^{\prime(f)}$ at $\zeta_2 = -1,\ P_c = -1$;

Figure 6: Components of the polarization vector of the scattered electron of its energy $\zeta_2^{\prime(f)}$, $\zeta_3^{\prime(f)}$, $\zeta_3^{\prime(f)}$ at $\zeta_1 = -1,\ P_c = -1$;



1.0
0.8
0.6
0.4
0.2
0.0
0.0
0.2
0.4
0.6
0.8
1.0
$\zeta_3'^{f}$
Y_1



1.0
0.8
0.6
0.4
0.2
0.0
$\zeta_2'^f$
Y_1
0.0
0.2
0.4
0.6
0.8
1.0


1.0
0.5
0.0
-0.5
-1.0
ζ'^{f}_{3}
Y_1
0.0
0.2
0.4
0.6
0.8
1.0


0.0
-0.2
-0.4
-0.6
-0.8
-1.0
0.0
0.2
0.4
0.6
0.8
1.0
$\zeta_2'^{f}$
Y_1


0.8
0.4
0.0
-0.4
-0.8
-1.2
0.0
0.2
0.4
0.6
0.8
1.0
$\zeta_3'^f$
Y_1


0.0
-0.2
-0.4
-0.6
-0.8
-1.0
0.0
0.2
0.4
0.6
0.8
1.0
$\zeta_2^{'f}$
Y_1


0.4
0.0
-0.4
-0.8
-1.2
0.0
0.2
0.4
0.6
0.8
1.0
$\zeta_3'^f$
Y_1


0.0
-0.2
-0.4
-0.6
-0.8
-1.0
0.0
0.2
0.4
0.6
0.8
1.0
$\zeta_2^{'f}$
Y_1


$\zeta_3'^f$
Y_1
1.2
0.8
0.4
0.0
-0.4
-0.8
0.0
0.2
0.4
0.6
0.8
1.0


1.0
0.5
0.0
-0.5
-1.0
0.0
0.2
0.4
0.6
0.8
1.0
$\zeta_2^{'f}$
Y_1



$\zeta_3^{'f}$
Y_1
1.2
0.8
0.4
0.0
-0.4
-0.8
0.0
0.2
0.4
0.6
0.8
1.0


1.0
0.5
0.0
-0.5
-1.0
0.0
0.2
0.4
0.6
0.8
1.0
$\zeta_2^{'f}$
$\zeta_1^{'f}$
Y_1